

# Argentina

Capital City : "Buenos Aires"  
Official Language: "Spanish"  
Monetary Unit: "peso (ARS)"

## General Information



### TRAVEL RESTRICTIONS ARE IN PLACE FOR THIS COUNTRY

There are entry requirements for anyone who plans to travel to England from this country. [Please check here for further information](#). Advice may vary in [Scotland](#), [Wales](#), and [Northern Ireland](#).

The information on these pages should be used to research health risks and to inform the pre-travel consultation.

Due to COVID-19, travel advice is subject to rapid change. Countries may change entry requirements and close their borders at very short notice. Travellers must ensure they check current [Foreign, Commonwealth & Development Office \(FCDO\) travel advice](#) in addition to the [FCDO specific country page](#) (where available) which provides additional information on travel restrictions and entry requirements in addition to safety and security advice.

Travellers should ideally arrange an appointment with their health professional at least four to six weeks before travel. However, even if time is short, an appointment is still worthwhile. This appointment provides an opportunity to assess health risks taking into account a number of factors including destination, medical history, and planned activities. For those with pre-existing health problems, an earlier appointment is recommended.

All travellers should ensure they have [adequate travel health insurance](#).

A list of useful resources including advice on how to reduce the risk of certain health problems is available below.

## Resources

- [Food and water hygiene](#)
- [Insect and tick bite avoidance](#)
- [Personal safety](#)
- [Sexually transmitted infections](#)
- [Sun protection](#)

## Vaccine Recommendations

Details of vaccination recommendations and requirements are provided below.

## All Travellers

Travellers should be up to date with routine vaccination courses and boosters as [recommended in the UK](#). These vaccinations include for example [measles-mumps-rubella \(MMR\)](#) vaccine and diphtheria-tetanus-polio vaccine.

Country specific diphtheria recommendations are not provided here. Diphtheria tetanus and polio are combined in a single vaccine in the UK. Therefore, when a tetanus booster is recommended for travellers, diphtheria vaccine is also given. Should there be an outbreak of diphtheria in a country, diphtheria vaccination guidance will be provided.

Those who may be at increased risk of an infectious disease due to their work, lifestyle choice, or certain underlying health problems should be up to date with additional recommended vaccines. See the individual chapters of the 'Green Book' [Immunisation against infectious disease](#) for further details.

## Certificate Requirements

- There are no certificate requirements under International Health Regulations.
- There is a **risk of yellow fever** transmission in parts of Argentina (see '*Some Travellers*' section below).

## Most Travellers

The vaccines in this section are recommended for most travellers visiting this country. Information on these vaccines can be found by clicking on the blue arrow. Vaccines are listed alphabetically.

### Tetanus

Tetanus is caused by a toxin released from *Clostridium tetani* and occurs worldwide. Tetanus bacteria are present in soil and manure and may be introduced through open wounds such as a puncture wound, burn or scratch.

### Prevention

Travellers should thoroughly clean all wounds and seek appropriate medical attention.

### Tetanus vaccination

- Travellers should have completed a primary vaccination course according to the UK schedule.
- If travelling to a country where medical facilities may be limited, a booster dose of a tetanus-containing vaccine is recommended if the last dose was more than ten years ago even if five doses of vaccine have been given previously.

Country specific information on medical facilities may be found in the 'health' section of the [FCDO foreign travel advice](#) website.

[Tetanus in brief](#)

## Some Travellers

The vaccines in this section are recommended for some travellers visiting this country. Information on when these vaccines should be considered can be found by clicking on the arrow. Vaccines are listed alphabetically.

## Hepatitis A

Hepatitis A is a viral infection transmitted through contaminated food and water or by direct contact with an infectious person. Symptoms are often mild or absent in young children, but the disease becomes more serious with advancing age. Recovery can vary from weeks to months. Following hepatitis A illness immunity is lifelong.

Those at increased risk include travellers visiting friends and relatives, long-stay travellers, and those visiting areas of poor sanitation.

### Prevention

All travellers should take care with personal, food and water hygiene.

### Hepatitis A vaccination

Vaccination is recommended for those whose activities put them at increased risk. This includes:

- those who are staying with or visiting the local population.
- frequent and/or long-stay travellers to areas where sanitation and food hygiene are likely to be poor.
- adventure travellers visiting rural areas and staying in basic accommodation such as backpackers.
- those with existing medical conditions such as liver disease or haemophilia.
- men who have sex with men.
- injecting drug users.
- those who may be exposed to the virus through their work.
- those going to areas of hepatitis A outbreaks who have limited access to safe water and medical care.

[Hepatitis A in brief](#)

## Hepatitis B

Hepatitis B is a viral infection; it is transmitted by exposure to infected blood or body fluids. This mostly occurs during sexual contact or as a result of blood-to-blood contact (for example from contaminated equipment during medical and dental procedures, tattooing or body piercing procedures, and sharing of intravenous needles). Mothers with the virus can also transmit the infection to their baby during childbirth.

### Hepatitis B in Argentina

2% or more of the population are known or thought to be persistently infected with the hepatitis B virus (intermediate/high prevalence).

## Prevention

Travellers should avoid contact with blood or body fluids. This includes:

- avoiding unprotected sexual intercourse.
- avoiding tattooing, piercing, public shaving, and acupuncture (unless sterile equipment is used).
- not sharing needles or other injection equipment.
- following universal precautions if working in a medical/dental/high risk setting.

A sterile medical equipment kit may be helpful when travelling to resource poor areas.

## Hepatitis B vaccination

Vaccination could be considered for all travellers, and is recommended for those whose activities or medical history put them at increased risk including:

- those who may have unprotected sex.
- those who may be exposed to contaminated needles through injecting drug use.
- those who may be exposed to blood or body fluids through their work (e.g. health workers).
- those who may be exposed to contaminated needles as a result of having medical or dental care e.g. those with pre-existing medical conditions and those travelling for medical care abroad including those intending to receive renal dialysis overseas.
- long-stay travellers.
- those who are participating in contact sports.
- families adopting children from this country.

[Hepatitis B in brief](#)

## Rabies

Rabies is a viral infection which is usually transmitted following contact with the saliva of an infected animal most often via a bite, scratch or lick to an open wound or mucous membrane (such as on the eye, nose or mouth). Although many different animals can transmit the virus, most cases follow a bite or scratch from an infected dog. In some parts of the world, bats are an important source of infection.

Rabies symptoms can take some time to develop, but when they do, the condition is almost always fatal.

The risk of exposure is increased by certain activities and length of stay (see below). Children are at increased risk as they are less likely to avoid contact with animals and to report a bite, scratch or lick.

## Rabies in Argentina

Rabies is considered a risk and has been reported in domestic animals in this country. Bats may also carry rabies-like viruses.

### Prevention

- Travellers should avoid contact with all animals. Rabies is preventable with prompt post-exposure treatment.
- Following a possible exposure, wounds should be thoroughly cleansed and an urgent local medical assessment sought, even if the wound appears trivial.
- Post-exposure treatment and advice should be in accordance with [national guidelines](#).

### Rabies vaccination

A full course of pre-exposure vaccines simplifies and shortens the course of post-exposure treatment and removes the need for rabies immunoglobulin which is in short supply world-wide. Pre-exposure vaccinations are recommended for travellers whose activities put them at increased risk including:

- those at risk due to their work (e.g. laboratory staff working with the virus, those working with animals or health workers who may be caring for infected patients).
- those travelling to areas where access to post-exposure treatment and medical care is limited.
- those planning higher risk activities such as running or cycling.
- long-stay travellers (more than one month).

[Rabies in brief](#)

### Typhoid

Typhoid is a bacterial infection transmitted through contaminated food and water. Previous typhoid illness may only partially protect against re-infection.

Travellers who will have access to safe food and water are likely to be at low risk. Those at increased risk include travellers visiting friends and relatives, frequent or long-stay travellers to areas where sanitation and food hygiene are likely to be poor, and laboratory personnel who may handle the bacteria for their work.

### Typhoid in Argentina

Typhoid fever is known or presumed to occur in this country.

### Prevention

All travellers should take care with personal, food and water hygiene.

### Typhoid vaccination

- Vaccination could be considered for those whose activities put them at increased risk (see

above).

- Oral and injectable typhoid vaccinations are available.

[Typhoid in brief](#)

## Yellow Fever

Yellow fever is a viral infection transmitted by mosquitoes which predominantly feed between dawn and dusk, but may also bite at night, especially in the jungle environment. Symptoms may be absent or mild, but in severe cases it can cause internal bleeding, organ failure and death.

### Yellow fever in Argentina

There is a risk of yellow fever transmission in **parts of this country** (see below).

#### Prevention

Travellers should avoid mosquito bites at all times.

#### Yellow fever vaccination

- Vaccination is recommended for travellers aged 9 months and older travelling to Corrientes and Misiones Provinces including the Iguazú Falls.
- Vaccination is **generally not recommended** for travel to areas with low potential for exposure to yellow fever: Formosa Province and designated areas of Chaco, Jujuy and Salta Provinces (see map).
- People aged 60 years or older **should not** be given the vaccine for travel to these areas with low potential for exposure due to a higher risk of life-threatening side effects.
- Vaccination could be considered for a small subset of travellers (aged 9 months to less than 60 years of age) for travel to areas with low potential for exposure to yellow fever who are at increased risk because of:
  - Prolonged travel.
  - Heavy exposure to mosquitoes.
  - Inability to avoid mosquito bites.
- Vaccination is **not recommended** for travellers whose itineraries are limited to areas and provinces not listed above.
- See vaccine recommendation map below.

The yellow fever vaccine is not suitable for all travellers, there are specific undesirable effects associated with it. This vaccine is only available at [registered yellow fever vaccination centres](#). Health professionals should carefully assess the risks and benefits of the vaccine, and seek specialist advice if necessary.

[Yellow fever in brief](#)

## Yellow fever vaccine recommendations in Argentina

[Map provided by the Travelers' Health Branch, Centers for Disease Control and Prevention](#)







**Current as of August 2018. This map, which aligns with recommendations also published by the World Health Organization (WHO), is an updated version of the 2010 map created by the Informal WHO Working Group on the Geographic Risk of Yellow Fever.**

**2. In 2017-8, the yellow fever vaccination recommendations for travellers to Brazil were expanded due to a large outbreak of yellow fever in multiple states within that country.**

**3. Yellow fever (YF) vaccination is generally not recommended in areas where there is low potential for YF virus exposure. However, vaccination might be considered for a small subset of travelers to these areas who are at increased risk for exposure to YF virus because of prolonged travel, heavy exposure to mosquitoes, or inability to avoid mosquito bites. Consideration for vaccination of any traveler must take into account the traveler's risk of being infected with YF virus, country entry requirements, and individual risk factors for serious vaccine-associated adverse events (e.g. age, immune status).**

## **Malaria**

- There is no risk of malaria in Argentina.
- Bite avoidance is recommended in all areas.

## **Other Risks**

There are some risks that are relevant to all travellers regardless of destination. These may for example include road traffic and other accidents, diseases transmitted by insects or ticks, diseases transmitted by contaminated food and water, sexually transmitted infections, or health issues related to the heat or cold.

Some additional risks (which may be present in all or part of this country) are mentioned below and are presented alphabetically. Select risk to expand information.

### **Altitude**

There is a risk of altitude illness when travelling to destinations of 2,500 metres (8,200 feet) or higher. Important risk factors are the altitude gained, rate of ascent and sleeping altitude. Rapid ascent without a period of acclimatisation puts a traveller at higher risk.

There are three syndromes; acute mountain sickness (AMS), high-altitude cerebral oedema (HACE) and high-altitude pulmonary oedema (HAPE). HACE and HAPE require immediate descent and medical treatment.

#### **Altitude illness in Argentina**

There is a point of elevation in this country higher than 2,500 metres. An example place of interest: Aconcagua 6,960m.

#### **Prevention**

- Travellers should spend a few days at an altitude below 3,000m.
- Where possible travellers should avoid travel from altitudes less than 1,200m to altitudes greater than 3,500m in a single day.

- Ascent above 3,000m should be gradual. Travellers should avoid increasing sleeping elevation by more than 500m per day and ensure a rest day (at the same altitude) every three or four days.
- Acetazolamide can be used to assist with acclimatisation, but should not replace gradual ascent.
- Travellers who develop symptoms of AMS (headache, fatigue, loss of appetite, nausea and sleep disturbance) should avoid further ascent. In the absence of improvement or with progression of symptoms the first response should be to descend.
- Development of HACE or HAPE symptoms requires immediate descent and emergency medical treatment.

[Altitude illness in brief](#)

### Biting insects or ticks

Insect or tick bites can cause irritation and infections of the skin at the site of a bite. They can also spread certain diseases.

#### Diseases in South America

There is a risk of insect or tick-borne diseases in some areas of South America. This includes diseases such as [American Trypanosomiasis \(Chagas Disease\)](#), [chikungunya](#), [leishmaniasis](#) and [West Nile virus](#).

#### Prevention

- All travellers should avoid insect and tick bites day and night.
- There are no vaccinations (or medications) to prevent these diseases.

Further information about specific insect or tick-borne diseases for this country can be found, if appropriate on this page, in other sections of the country information pages and the [insect and tick bite avoidance factsheet](#).

### Dengue

Dengue is a viral infection transmitted by mosquitoes which predominantly feed between dawn and dusk. It causes a flu-like illness, which can occasionally develop into a more serious life-threatening form of the disease. Severe dengue is rare in travellers.

The mosquitoes that transmit dengue are most abundant in towns, cities and surrounding areas. All travellers to dengue areas are at risk.

#### Dengue in Argentina

There is a risk of dengue in the provinces north of and including the province of Buenos Aires. Affected provinces include: Buenos Aires, Buenos Aires D.F., Catamarca, Chaco, Cordoba, Corrientes, Entre Rios, Formosa, Jujuy, Misiones, Salta, Santa Fe, Santiago Del Estero, and Tucuman.

## Prevention

- All travellers should avoid mosquito bites particularly between dawn and dusk.
- There is no vaccination or medication to prevent dengue.

[Dengue in brief](#)

## Influenza

Seasonal influenza is a viral infection of the respiratory tract and spreads easily from person to person via respiratory droplets when coughing and sneezing. Symptoms appear rapidly and include fever, muscle aches, headache, malaise (feeling unwell), cough, sore throat and a runny nose. In healthy individuals, symptoms improve without treatment within two to seven days. Severe illness is more common in those aged 65 years or over, those under 2 years of age, or those who have underlying medical conditions that increase their risk for complications of influenza.

### Seasonal influenza in Argentina

Seasonal influenza occurs throughout the world. In the northern hemisphere (including the UK), most influenza occurs from as early as October through to March. In the southern hemisphere, influenza mostly occurs between April and September. In the tropics, influenza can occur throughout the year.

## Prevention

All travellers should:

- Avoid close contact with symptomatic individuals
- Avoid crowded conditions where possible
- Wash their hands frequently
- Practise 'cough hygiene': sneezing or coughing into a tissue and promptly discarding it safely, and washing their hands
- Avoid travel if unwell with influenza-like symptoms
- A vaccine is available in certain circumstances (see below)\*

**\*In the UK, seasonal influenza vaccine is offered routinely each year to those at higher risk of developing of severe disease following influenza infection, and certain additional groups such as healthcare workers and children as part of the UK national schedule (see [information on vaccination](#)). For those who do not fall into these groups, vaccination may be available privately.**

If individuals at higher risk of severe disease following influenza infection are travelling to a country when influenza is likely to be circulating they should ensure they received a flu vaccination in the previous 12 months.

The vaccine used in the UK protects against the strains predicted to occur during the winter months of the northern hemisphere. It is not possible to obtain vaccine for the southern hemisphere in the

UK, but the vaccine used during the UK influenza season should still provide important protection against strains likely to occur during the southern hemisphere influenza season, and in the tropics.

### **Avian influenza**

Avian influenza viruses can rarely infect and cause disease in humans. Such cases are usually associated with close exposure to infected bird or animal populations. Where appropriate, information on these will be available in the outbreaks and news sections of the relevant country pages. Seasonal influenza vaccines will not provide protection against avian influenza.

[Avian influenza in brief](#)

### **Outdoor air quality**

Poor air quality is a significant public health problem in many parts of the world. Exposure to high levels of air pollution over short time periods (e.g. minutes/hours/days) and longer time periods (e.g. years) is linked to many different acute and chronic health problems. These effects are mainly on the respiratory (lungs and airways) and cardiovascular (heart function and blood circulation) systems.

Current information on world air quality is available from the [world air quality index project](#).

### **Prevention**

Travellers with health problems that might make them more vulnerable to the effects of air pollution who are travelling to areas of high pollution should:

- discuss their travel plans with their doctor, and carry adequate supplies of their regular medication
- take sensible precautions to minimise their exposure to high levels of air pollution
- check local air quality data and amend their activities accordingly
- take notice of any health advisories published by the local Ministry of Health and Department for Environment, and follow the guidance provided.

It is unclear if face masks are beneficial at reducing exposure and may make breathing more difficult for those with pre-existing lung conditions. Those who choose to use one should make sure that the mask fits well and know how to wear it properly.

[Outdoor air quality in brief](#)

### **Zika Virus**

Zika virus (ZIKV) is a viral infection transmitted by mosquitoes which predominantly feed between dawn and dusk. A small number of cases of sexual transmission of ZIKV have also been reported. Most people infected with ZIKV have no symptoms. When symptoms do occur they are usually mild and short-lived. Serious complications and deaths are not common. However ZIKV is a cause of Congenital Zika Syndrome (microcephaly and other congenital anomalies) and neurological complications such as Guillain-Barré syndrome.

### **Zika virus in Argentina**

There is a risk of ZIKV in this country. **Pregnant women should consider avoiding travel** to this country until after the pregnancy. In the event that travel is unavoidable, the pregnant traveller must be informed of the risks which ZIKV presents.

Details of specific affected areas within this country are limited; confirmed cases are reported from the provinces of Chaco, Formosa, Salta, and Tucumán. The mosquitoes that transmit ZIKV are unlikely to be found above 2,000m altitude. There are a number of areas above 2,000m in the Andes mountains.

The map below shows areas which are above 2,000m and can be used by travellers and health professionals as a general guide to indicate potentially lower risk areas for mosquito-acquired ZIKV infection. Travellers whose itineraries are limited to areas above 2,000m are at a lower risk of acquiring ZIKV from a mosquito; however there may still be a risk of sexual transmission.

[Map provided by the Travelers' Health Branch, Centers for Disease Control and Prevention](#)



***The categories shown on this map are intended as a general guideline and should not be considered to indicate absolute risk. Elevation may vary within an area to a larger extent than this map can depict. The presence of mosquitoes may change seasonally, with increasing temperatures or rainfall, and may change over time. Travellers to destinations that cross or are near an elevation border may wish to consider the destination as an area of lower elevation. Travellers to high elevations are still at risk of getting Zika from sex.***

## Prevention

- All travellers should avoid mosquito bites particularly between dawn and dusk.
- There is no vaccination or medication to prevent ZIKV infection.
- Women should avoid becoming pregnant while travelling in this country, and for 2 months (8 weeks) after their last possible ZIKV exposure\* (see below if male partner has travelled).
- If a woman develops symptoms compatible with ZIKV infection, it is recommended she avoids becoming pregnant for a further 2 months following recovery.
- Pregnant women who visited this country while pregnant, or who become pregnant within 2 months after their last possible ZIKV exposure\*, should contact their GP, obstetrician or midwife for further advice, even if they have not been unwell.

Please note screening of returning travellers without ZIKV symptoms is not available on the NHS. Couples planning pregnancy in the very near future should consider whether they should avoid travel to a country or area with risk of ZIKV, rather than delay conception for the recommended period (see below) after travel. This particularly includes couples in assisted fertility programmes.

### Preventing sexual transmission

Couples should follow [guidance on prevention of sexual transmission of ZIKV](#) and avoid conception as follows:

- If both partners travelled, for 3 months after last possible ZIKV exposure\*
- Male traveller only, for 3 months after last possible ZIKV exposure\*
- Female traveller only, for 2 months after last possible ZIKV exposure\*

See [further information for pregnant women, their partners and couples planning pregnancy](#).

**\*Last possible Zika virus exposure is defined as the later of either the date of leaving a country or area with risk for ZIKV transmission, or the date on which unprotected sexual contact with a potentially infectious partner took place.**

See [detailed guidance on factors to consider when assessing the risk of ZIKV](#).

[Zika virus in brief](#)

## Covid-19

COVID-19 disease is caused by the coronavirus SARS-CoV2. The main symptoms of COVID-19 are recent onset of a new continuous cough and/or a high temperature and/or a loss of, or change in, normal sense of taste or smell. Symptoms range from mild to life-threatening. Older people and those with underlying health problems are more likely to develop severe disease. Current evidence also shows that there is a higher risk of infection and of severe disease in Black, Asian and Minority Ethnic groups in the UK. The reasons for this are currently not clear.

The virus is mainly transmitted from human to human by breathing in droplets from the nose or mouth of an infected person or by touching the infected droplets on surfaces, then touching the eyes, nose or mouth. A COVID-19 vaccine is now available in some countries, but even if

vaccinated, prevention still relies on strict attention to social distancing guidance and personal and respiratory hygiene.

Travellers should always check the UK [Foreign, Commonwealth & Development Office \(FCDO\) travel advice](#) and their [country-specific pages](#) for the latest COVID-19 travel advisories which may include information on travel restrictions, quarantine or COVID-19 testing requirements. This includes considering the recommendations and requirements for any transit countries. Travellers should be aware that case numbers in individual countries/areas can increase rapidly and country requirements can change at short notice. They should check this page and FCDO travel advice regularly for updates.

## Covid-19 In Argentina

Based on currently available information, there is a high risk of exposure to COVID-19 in this country. Due to the emerging COVID-19 variant in this region/country, all travellers are advised to avoid non-essential travel to this country.

There are entry requirements for anyone who plans to travel to England from this country. [Please check here for further information](#). Advice may vary in [Scotland](#), [Wales](#), and [Northern Ireland](#).

## Prevention

If travelling to this country travellers should:

- Discuss their plans with a health professional. This is particularly important for those at higher risk of severe COVID-19.
- Consider the risk at all destinations including any transit countries and the risk during travel itself.
- Check with the airline/tour operator about physical distancing measures or other measures in place to reduce risk during travel.
- Follow the latest social distancing guidance, including any local requirements and maintain good hand, respiratory, and personal hygiene at all times. This may be particularly important if staying with friends and family.
- Complete the full COVID-19 vaccination schedule when offered this as part of the UK vaccination programme.

See [guidance on factors to consider when assessing the risk of COVID-19 for travellers](#).

If travellers develop COVID-19 symptoms while abroad, they should:

- Follow local guidelines on self-isolation, testing and avoiding travel.
- Contact their travel insurance provider.
- Seek medical advice if needed.

[COVID-19 in brief](#)

[Latest News](#)

[Latest Outbreaks](#)

---