

Travelling to visit friends and relatives

VFR is the second most common reason given for travelling overseas from the UK

Key messages

- Travellers visiting friends and relatives (VFR) are a diverse group with a wide variation in country of origin, immunisation status and travel destinations.
- VFR travellers are less likely to seek pre-travel advice. Even if time is short before departure, a last-minute appointment is still worthwhile.
- There is an increased risk of a range of preventable infections for some VFR travellers, including vaccine-preventable diseases and malaria.
- UK travellers visiting friends and relatives in Africa are at highest risk of getting malaria.
- VFR travellers are less likely to take malaria prevention tablets than other travellers to Africa. The importance of malaria prevention tablets in this group should be emphasised.

Overview

Visiting friends and relatives (VFR) is now the second most common reason given for travelling overseas from the UK, after travelling for a tourist holiday [1]. In this factsheet, a VFR traveller is defined as a person, ethnically and racially distinct from the majority population in the UK, who returns to his or her home country (lower-income country) to visit friends or relatives. Family members, such as the partner or children, who may have been born in the UK, are also included in this definition [2]. VFR travellers may include immigrants, asylum seekers/refugees and students.

VFR travellers have a greater risk of travel related diseases and other health risks [3, 4] and are less likely to seek pre-travel advice [5, 6]. A number of reasons may explain the disproportionate risk for this group of travellers: they may travel to resource-poor areas, have closer contact with the local population and travel for longer periods [7]. Barriers to VFR travellers receiving travel health advice may include factors such as; access to travel health services, cost of vaccines and malaria



chemoprophylaxis, perception of travel-related risks by the health care advisor and the traveller, and communication issues [4].

Pre-travel preparation

All travellers should review the country specific information including the <u>Foreign, Commonwealth & Development Office (FCDO)</u> advice on the <u>Country Information pages</u> and make an appointment with their GP practice or a travel clinic. Ideally travellers should seek advice at least four to six weeks before departure.

For individuals with complex medical problems an earlier appointment is recommended. However, it is recognised that VFR travellers often need to travel at short notice for various reasons such as a family emergency. If time is short, an appointment is still worthwhile.

Individuals with complex medical problems, older travellers, pregnant women, infants or young children often prove challenging for the health professional [6], especially if they present with little time prior to departure [6].

Pre-travel advice should be tailored to prioritise what is achievable within a short time frame.

Pre-travel advice should be given sensitively. Focussing on behaviours to reduce exposure to diseases, such as the use of mosquito nets and insect repellents and food/water hygiene may be more effective than emphasising the risks in the destination country [8].

Travellers are recommended to take out <u>travel insurance</u> to cover the potentially high cost of medical treatment and repatriation, if they are injured or fall ill abroad.

Food and water risks

Contaminated food and water can transmit many different infectious diseases; the risk is higher in low-income regions of the world [9]. VFR travellers are at increased risk of certain infections, for example typhoid. In 2023, of the 644 imported cases of typhoid/paratyphoid in England, Wales and Northern Ireland, where reason for travel from UK was known (n=546) 87 percent were reported in VFR travellers [10].

Care should be taken with <u>food and water hygiene</u>, and VFR travellers should know how to manage the symptoms of <u>travellers' diarrhoea</u> and when to seek medical advice. The effects of diarrhoea are generally greater in the very young, elderly and frail travellers. Those with special needs or certain medical problems should also take particular care for example those with a weak immune system, inflammatory bowel disease, chronic (long term) kidney or heart disease, pregnant women and those taking medication to reduce stomach acid [11].

It can be difficult for VFR travellers to follow all the recommended advice as they will not want to cause offence by refusing certain drinks or food. They may want to think about ways to get around



this such as emphasising that children born in the UK won't have immunity to certain diseases, mention that they are following doctors' instructions or have a sensitivity to certain foods. Having hand sanitising gel or wipes ready to use discreetly before eating may also be helpful.

Vector-borne risks

In addition to malaria, there are many <u>diseases spread by mosquitoes and other insects (vectors)</u>. For many of these there are no vaccines or medications to prevent infection and so <u>insect bite avoidance measures</u> are vital.

Malaria

Malaria is a serious, potentially life-threatening infection spread by mosquitoes. The World Health Organization estimated 263 million cases of malaria occurred worldwide in 2023 [12], with 94 percent of the cases in the African region. Studies on the knowledge and perception of malaria among VFR travellers in the UK have shown that the majority are aware of the potential risk of contracting the disease. However, malaria is commonly not regarded as a serious health threat amongst this group [13].

In the UK, malaria mainly affects the non-UK born population and their families, particularly those from Africa and South Asia. Those visiting friends and relatives in Africa are at highest risk of acquiring malaria abroad [14].

Travellers of African origin may overestimate the amount of protection that having been brought up in Africa may give them. Those who are born in a malarious area often suffer repeated infections during early childhood which can bring about a state of partial immunity. However, past infection does not confer long-lasting immunity, and any immunity acquired rapidly disappears if the individual moves to an area where malaria does not occur [14].

There is no such thing as "natural" immunity to malaria. Second-generation members of VFR families who migrate to the UK will not have any immunity to malaria, and their children will be vulnerable.

Data suggests that VFR travellers are significantly less likely to take malaria prevention tablets than other travellers to Africa, and are less likely to seek pre-travel malaria prevention advice [14]. Possible reasons for this include health beliefs, such as believing malaria is a mild disease, or that having sickle cell disease affords immunity to malaria.

VFR travellers may be under pressure not to use malaria prevention tablets or bite prevention methods as it is thought to be unnecessary, or they may not want to feel an 'outsider' in their community of origin.

The cost of malaria prevention tablets may also deter VFR travellers, especially if travelling as a family. The importance of malaria prevention tablets should be emphasised. They should always be



purchased from a reputable source in the UK, started before travel and continued as directed for maximum protection.

All travellers should be aware of the signs and symptoms of malaria and should be advised to seek immediate urgent medical attention if these occur either whilst abroad or up to a year after their return. They should also let the health professional know they have visited a malaria risk area.

COVID-19

All individuals should follow <u>current UK recommendations</u> to reduce their risk of catching COVID-19 and passing it on to others.

Some people are at increased risk of severe infection from COVID-19. This mainly relates to age and any existing medical conditions and includes those prioritised for COVID-19 vaccination (see Chapter 14a; Table 2 and 3 in Immunisation against Infectious Disease).

Travellers who are at increased risk of severe COVID-19 disease should assess their individual circumstances, including medical facilities at their destination, and consider whether postponing travel would be appropriate.

General guidance regarding <u>risk assessment for travel</u> during the COVID-19 pandemic and information about the <u>COVID-19 vaccination programme</u> is available.

Vaccination

Travellers should be up to date with routine vaccination courses and boosters as recommended in the <u>UK schedule</u>. Information on primary vaccinations provided overseas can be found on the <u>World Health Organization website</u> and guidance on <u>catching up individuals with uncertain or incomplete immunisation status</u> is available from the UK Health Security Agency.

There is evidence that those visiting friends and family are more likely to acquire vaccine preventable diseases than tourists [15]. VFR travellers may be more likely to decline pre-travel vaccinations, one study has shown that of vaccine refusers, 38% were VFR travellers [7].

Pregnancy

<u>Pregnancy</u> can increase the risks from travel-related diseases, diseases may have a more severe progression, and some methods of disease prevention or treatment may be 'off licence', i.e. not covered on the manufacturers marketing authorisation. Generally, the use of inactivated vaccines during pregnancy is considered safe, but the recommendations should be based on a risk-and-benefit assessment [16]. Further information on vaccinations can be found in our <u>Pregnancy factsheet</u>.

Malaria infection during pregnancy is a significant public health problem with substantial risks for

the pregnant woman and her foetus. Pregnant women have an increased risk of severe malaria and fatality compared to non-pregnant women. Malaria can also cause anaemia, spontaneous abortion, stillbirth, prematurity or low birthweight. The UK Malaria Expert Advisory Group (UKMEAG) recommends that pregnant women should be advised against travel to malarious areas, particularly if there is chloroquine resistant malaria at the destination [14]. Pregnant travellers should be particularly vigilant with <u>insect bite avoidance measures</u> and <u>food and water hygiene</u>.

Children

<u>Infants/children</u> are at greater risk of a severe disease progression of travel related infections; e.g. malaria [14, 17]. The UK Malaria Expert Advisory Group recommends that parents are advised against taking infants and young children to malarious areas without adequate precautions [14].

Some vaccines are not recommended for infants or young children below a certain age e.g. <u>yellow fever</u> and <u>typhoid</u>. Young children can be at particular risk of exposure to <u>rabies</u>, as they may be more inclined to approach domestic dogs and cats and may not report an exposure; 40 percent of rabies cases are in children under 15 years of age [18].

Other health risks

Animal bites

VFR travellers may have a greater risk of <u>rabies</u>. Staying in relatives' homes may lead to increased exposure to domestic and wild animals compared to those staying in hotels and tourist resorts [19]. Also, VFR travellers may have been more likely to decline pre-exposure vaccination [19].

With the exception of Antarctica, <u>rabies is endemic on all continents</u>. Rabies occurs in warm-blooded mammals, commonly dogs, but also other domestic and wild animals including bats. The virus is transmitted to humans by a bite or scratch, or when saliva from an infected animal comes into contact with broken skin or mucous membranes (eyes, nose, or mouth). Rabies is almost invariably fatal once symptoms appear [18], but the disease is preventable if the correct post-exposure treatment (PET) is provided quickly. Accessing correct PET can be challenging in some countries.

Sexually transmitted infections

VFR travellers have been shown to be at greater risk of human immunodeficiency virus (HIV) and sexually transmitted infections compared with tourists and business travellers to the same destination [8]. Those travelling without a steady partner should be aware of the risks; reliable condoms should be carried and used to reduce the risks.

Female genital mutilation (FGM)

The <u>World Health Organization</u> estimates that more than 230 million girls and women alive today have undergone <u>Female genital mutilation (FGM)</u>. It is an offence for FGM to be carried out on a UK citizen or resident of the UK or for such a person to be taken abroad for this purpose [20]. Health professionals have a <u>mandatory reporting duty</u> if FGM is identified or a child is thought to be at risk. Failure to do so may result in <u>disciplinary procedures</u> through the health professional's regulatory body.

Medications

VFR travellers may be tempted to purchase medications, particularly malaria chemoprophylaxis at their destination. There are concerns about fake medicines in some parts of the world causing harm to patients and failing to treat/prevent the disease for which they were intended [21]. It is generally safer to obtain medication in the UK before travel. Additionally, a course of malaria chemoprophylaxis needs to be started prior to exposure to malaria [14], so purchasing it on arrival at the destination may mean that the course is started later than recommended. See our Medicines and travel factsheet for further details.

General advice for those who get sick abroad

Travellers with fever, diarrhoea with blood or any other worrying symptoms, such as altered mental status, severe abdominal pain, jaundice or rash should seek prompt medical care. Those who need to seek medical advice abroad should contact their insurance company and keep receipts so they can claim back the costs from the insurance or EHIC as appropriate.

Resources

- UK Health Security Agency: Travelling overseas to visit friends and relatives?
- Pregnancy
- Children

REFERENCES

- 1. Office for National Statistics: Travel Trends 2023. August 2024 [Accessed 17 December 2024]
- 2. Wanduragala D, Coyle C, Angelo K, Stauffer W. Visiting Friends and Relatives: VFR Travel. In US Centers for Disease Control (CDC) Yellow Book, Health Information for International Travel 2024 [Accessed 17 December 2024]
- **3.** Barnet E, MacPherson D, Stauffer W et al. The visiting friends or relatives traveler in the 21st century: time for a new definition. JTM 2010; volume 17(3): 163-170 [Accessed 3 September 2021]
- **4.** LaRocque R, Deshpande B, Rao S et al. Pre-travel health care of immigrants returning to home to visit friends and relatives. Am J Trop Med Hyg; 2013; 88(2): 376-380 [Accessed 3 September 2021]
- **5.** Warne B, Weld LH, Cramer JP, et al Travel Related Infection in European Travellers, EuroTravNet 2011 Journal Of Travel Medicine, 2014 21(4) 248-254 [Accessed 3 September 2021]
- **6.** Gurgle H, Rosel D, Erickson T et al. Impact of traveling to visit friends and relatives on chronic disease management. JTM 2013; 20(2): 95-100 [Accessed 3 September 2021]

- 7. Lammert S, Rao S, Jentes E et al. Refusal of recommended travel-related vaccines among U.S. international travellers in Global TravEpiNet J Travel Med. 2017 Jan: 24(1) [Accessed 3 September 2021]
- 8. Behrens R, Leder K. Chapter 31; Visiting friends and relatives. In Keystone J, Kozarsky P, Freedman D et al. Travel Medicine. 3rd Edition. 2013. Elsevier.
- 9. Steffen, R., Hill, D.R., DuPont, H.L. Traveler's diarrhea; a clinical review. JAMA. 2015; 313 (1): 71-80. [Accessed 3 September 2021]
- **10.** UK Health Security Agency. Enteric fever (typhoid and paratyphoid) England, Wales and Northern Ireland: 2023. Updated 22 August 2024 [Accessed 17 December 2024]
- **11.** Bavishi C, Dupont HL Systematic review: the use of proton pumps inhibitors and increased susceptibility to enteric infection, Aliment Pharmacol Ther. 2011 Dec; 34(11-12):1269-81 [Accessed 3 September 2021]
- 12. World Health Organization. World Malaria Report 2024
- **13.** Behrens R, Neave P, Jones C. Imported malaria among people who travel to visit friends and relatives: is current UK policy effective or does it need a strategic change? Malaria Journal 2015;14:149
- **14.** Chiodini PL, Patel D, Goodyer L, Frise C, Mabayoje D, Aggarwal D. Guidelines for malaria prevention in travellers from the United Kingdom, 2024. London: UK Health Security Agency; October 2024 [Accessed December 17 2024]
- **15.** Boggild A, Castelli F, Gautret P et al. Vaccine preventable diseases in returned international travelers: results from the GeoSentinel Surveillance Network Vaccine 2010; 28: 7389-7395 [Accessed 3 September 2021]
- **16.** Matteelli A, Carvalho A & Bigoni S. Visiting relatives and friends (VFR), pregnant and other vulnerable travelers. Infect. Dis Clin N Am (2012); 26: 625-635 [Accessed 3 September 2021]
- 17. World Health Organization. Factsheet; Malaria. 4 December 2023 [Accessed 11 March 2024]
- 18. World Health Organization. Rabies vaccines position paper April 2018
- 19. Dolan S, Jentes E, Sotir M et al. Pre-exposure rabies vaccination among US international travellers: Findings from the Global TravEpiNet Consortium. Vector-Borne and Zoonotic Disease 2014; February: Vol 14(2) [Accessed 3 September 2021]
- **20.** Office for Health Improvement and Disparities. Female genital mutilation (FGM); migrant health guide. Updated 13 September 2021 [Accessed 11 March 2024]
- **21.** World Health Organization. Substandard and falsified medical products. 3 December 2024 [Accessed 17 December 2024]

Published Date: 05 Dec 2019

Updated Date: 17 Dec 2024